

Management of Hip Fractures in the Elderly: Timing of Surgical Intervention

Performance Measure – Technical Report

Adopted by:

The American Academy of Orthopaedic Surgeons Board of Directors
April 14, 2018

Contents

Burden of Disease	3
Etiology.....	3
Incidence and Prevalence.....	3
Roster	5
Methods	6
Evidence Base	6
Hip Fractures: Timing of Surgical Intervention.....	8
Measure Specifications	8
Validity	9
Reliability.....	10
Performance Gap	11
References.....	14
Appendix A. Disclosures	15

Executive Summary

This report presents one quality measure related to management of hip fractures in the elderly.

- Hip fractures: Timing of Surgical Intervention

This report presents the measure specifications and analytic results. Included are the rationale for each measure and the specific proposed technical approach to each measure.

BURDEN OF DISEASE

The economic burden of managing elderly hip fractures was estimated at \$17-20 billion in 2010.^{M1, M2}

A typical patient with a hip fracture spends US \$40,000 in the first year following hip fracture for direct medical costs and almost \$5,000 in subsequent years.

Costs to be considered include:

1. Direct Medical Cost
2. Long-term Medical Cost
3. Home Modification Costs
4. Nursing Home Costs

ETIOLOGY

Hip fractures in the elderly are the result of low energy trauma and often are associated with osteoporosis/low bone mass and other associated medical conditions that may increase the prevalence of falls.

INCIDENCE AND PREVALENCE

There was an estimated 340,000 hip fracture patients per year in United States in 1996 with most fractures occurring in women older than age 65 years, and an annual worldwide incidence of approximately 1.7 million.^{M1, M7}

Between 1986 and 2005, the annual mean number of hip fractures was 957.3 per 100,000 (95% confidence interval [CI], 921.7-992.9) for women and 414.4 per 100 000 (95% CI, 401.6-427.3) for men.^{M1}

With rising life expectancy, the number of elderly individuals and those with chronic health conditions is increasing and it is estimated that the prevalence of hip fractures will continue to increase. The number of people older than age 65 years is expected to increase from 37.1 million to 77.2 million by the year 2040, and the occurrence of hip fractures is expected increase concomitantly, with an estimated 6.3 million hip fractures predicted worldwide by 2050.^{M7}

Work Group Composition

Solicitation of the work group members was done through medical societies and research organizations that have a relevant interest in the selected topic, commonly treat/interact with patients who receive the procedure, or have particular expertise in measure development. This project's workgroup consisted of seventeen individuals.

Roster

1. **Timothy Brox, MD - Oversight Chair**
2. **Steven Olson, MD – Chair**
Orthopaedic Trauma Association
3. **David B. Carmack, MD**
American Academy of Orthopaedic Surgeons
4. **Charles M. Davis III, MD**
American Academy of Orthopaedic Surgeons
5. **Eitan Dickman, MD**
American College of Emergency Physicians
6. **Daniel Ari Mendelson, MS, MD, FACP, AGSF, CMD**
American Geriatrics Society
7. **Anna Noel Miller, MD**
American Academy of Orthopaedic Surgeons
8. **Arvind D. Nana, MD**
American Academy of Orthopaedic Surgeons
9. **Laura Lowe Tosi, MD**
American Academy of Orthopaedic Surgeons/ U.S. Bone and Joint Initiative
10. **Stephen L. Kates, MD**
Orthopaedic Trauma Association/ American Orthopaedic Association
11. **James F. Kellam, MD**
American Academy of Orthopaedic Surgeons
12. **Douglas W. Lundy, MD**
Orthopaedic Trauma Association
13. **Kevin Means, MD**
American Academy of Physical Medicine and Rehabilitation
14. **Simon Mears, MD**
American Academy of Orthopaedic Surgeons
15. **Colleen Walsh, DNP, RN**
Representative of National Association of Orthopaedic Nurses
16. **Douglas M. White, DPT**
Representative of American Physical Therapy Association
17. **Douglas G. Wright, MD**
Representative of American Academy of Orthopaedic Surgeons

Methods

This measure was developed by following the American Academy of Orthopaedic Surgeons' performance measures methodology:

<https://www.aaos.org/uploadedFiles/PreProduction/Quality/Measures/PM%20Methodology%20Final%20-%20July%202017.pdf>





Evidence Base

The quality measure in this document is based directly on the American Academy of Orthopaedic Surgeons Management of Hip Fractures in the Elderly Clinical Practice Guideline, published in 2014.

Prior to performing the literature search for this guideline, both patients and payors were surveyed for topics of interest related to the management of hip fractures in the elderly. These responses helped inform the PICO development by the workgroup. All PICO questions and inclusion criteria were developed a priori. AAOS staff trained in research methodology conducted a comprehensive systematic literature review, and final recommendations were developed by a panel of experts. The workgroup that created these final recommendations is separate from the one that evaluated these quality measures. All included articles underwent study design quality appraisal, which assessed risks of bias/confounders that may skew the study's results. Only the best available evidence was considered for inclusion in recommendations.

Requirements for the strength of recommendation are listed below as Table 1.

Table 1. Strength of Recommendation Descriptions

Strength	Overall Strength of Evidence	Description of Evidence Quality	Strength Visual
Strong	Strong	Evidence from two or more "High" quality studies with consistent findings for recommending for or against the intervention.	
Moderate	Moderate	Evidence from two or more "Moderate" quality studies with consistent findings, or evidence from a single "High" quality study for recommending for or against the intervention.	
Limited	Low Strength Evidence or Conflicting Evidence	Evidence from one or more "Low" quality studies with consistent findings or evidence from a single "Moderate" quality study recommending for against the intervention or diagnostic or the evidence is insufficient or conflicting and does not allow a recommendation for or against the intervention.	
Consensus*	No Evidence	There is no supporting evidence. In the absence of reliable evidence, the guideline work group is making a recommendation based on their clinical opinion. Consensus statements are published in a separate, complimentary document.	

Detailed methodology for guideline development can be found at:
https://www.aaos.org/uploadedFiles/PreProduction/Quality/Guidelines_and_Reviews/guidelines/Guideline%20and%20Systematic%20Review%20Processes_v2.0_Final.pdf.

Hip Fractures: Timing of Surgical Intervention

Measure Specifications

2018 Options for Individual Measures:

Claims, EMR, Registry

Measure Type:

Process

Description:

Percentage of patients (65 years and older) who present to the emergency department with a hip fracture receive surgical intervention within 48 hours of admission to the hospital.

Instructions:

This measure is to be reported at **each denominator eligible visit** occurring during the reporting period for patients age 65 or older admitted to the hospital with a low energy hip fracture during the reporting period. This measure may be reported by eligible clinicians who perform the quality actions described in the measure based on the services provided and the measure-specific denominator coding.

Measure Reporting:

The listed denominator criteria are used to identify the intended patient population. The numerator quality-data codes included in this specification are used to submit the quality actions allowed by the measure. All measure-specific coding should be reported on the claim(s) representing the eligible encounter. This measure is designed to be reported at the group/practice, hospital/facility/agency, or regional level.

Denominator:

Number of patients age 65 or older admitted to the hospital with a low energy hip fracture

Denominator Criteria (Eligible Cases):

ICD-10-CM: S72.00, S72.001, S72.002, S72.009, S72.01, S72.011, S72.012, S72.019, S72.02, S72.03, S72.032, S72.033, S72.034, S72.035, S72.036, S72.04, S72.041, S72.042, S72.043, S72.044, S72.045, S72.046, S72.05, S72.051, S72.052, S72.059, S72.060, S72.09, S72.091, S72.092, S72.099, S72.136, S72.14, S72.141, S72.143, S72.144, S72.145, S72.1346, S72.2, S72.21, S72.22, S72.23, S72.24, S72.25, S72.26

OR

ICD-9-CM: 820.8, 820, 820.02, 820.03, 820.09, 820.2, 820.21, 820.22

Numerator:

Number of patients in the denominator who are operated on within 48 hours of admission to the hospital.

Numerator Criteria (Eligible Cases):

CPT: 27235, 27236, 27244, 27245, 27248, 27254, 27269

Exclusions:

Patients that can be classified as having the following: non-operative fractures, multiple injuries, periprosthetic fracture, high energy trauma, and or meet local criteria for multiple trauma designation

Evidence-Based Recommendation:

Moderate evidence supports that hip fracture surgery within 48 hours of admission is associated with better outcomes.

Rationale

Nine moderate strength studies evaluated patient outcomes in relation to timing of hip fracture surgery (Elliot et al 25, Fox et al 26, McGuire et al 27, Moran et al 28, Novack et al 29, Orosz et al 30, Parker et al 31, Radcliff et al 32, Siegmeth et al 33). In many of these studies the presence of increased comorbidities represented a confounding effect, and therefore delays for medical reasons were often excluded.

The majority of studies favored improved outcomes in regards to mortality, pain, complications, or length of stay (Elliot et al 25, McGuire et al 27, Novack et al 29, Orosz et al 30, Parker et al 31, and Siegmeth et al 33). Although several studies showed a benefit of surgery within 48 hours, one study showed no harm with a delay up to four days for patients fit for surgery who were not delayed for medical reasons (Moran et al 28). Patients delayed due to medical reasons had the highest mortality and it is this subset of patients that could potentially benefit the most from earlier surgery.

Risks and Harms of Implementing this Recommendation

There are no known harms associated with implementing this recommendation.

Validity

Validity testing focuses on systematic errors and bias. It involves testing agreement between the data elements obtained when implementing the measure as specified and data from another source of known accuracy. Validity of computed measure scores involves testing hypotheses of relationships between the computed measure scores as specified and other known measures of quality or conceptually related aspects of quality. A variety of approaches can provide some evidence for validity. The specific terms and definitions used for validity may vary by discipline, including face, content, construct, criterion, concurrent, predictive, convergent, or discriminant validity.

The validity of this measure comes from a combination of a strong basis in literature, by way of a systematic literature review conducted for a clinical practice guideline, and the face validity vote of a panel of experts. This independent clinician group consisted of the Evidence-based Quality, Value, and Safety Committee from the Orthopaedic Trauma Association, a distinct entity from the American Academy of Orthopaedic Surgeons. The committee was provided with the measure description, specifications, evidence-base, and included data elements, and asked the direct question: “Could a low computed performance measure score (i.e. time to surgery consistently exceeding 48 hours) be used to identify poor care and subsequently distinguish between appropriate and poor-quality care?” This committee affirmed face validity based on the provided materials.

Clinical Data

The Measure was specified to include all patients who present to the ED with a low energy hip fracture requiring operative care. The population is derived from the most recent year of the 5% carrier files from the 5 years of the patient files from The Cleveland Clinic. The Cleveland Clinic Data represents all the Hip Fracture Patients from 2012 through 2016 (Table 2).

Table 2. Description of Cleveland Clinic Data Files

Year	Cleveland Clinic Data	Size
2012	Hip Fracture Patients	N=447
2013	Hip Fracture Patients	N=469
2014	Hip Fracture Patients	N=425
2015	Hip Fracture Patients	N=470
2016	Hip Fracture Patients	N=528

Reliability

Reliability was calculated according to the methods outlined in a technical report prepared by J.L. Adams titled “The Reliability of Provider Profiling: A Tutorial” (RAND Corporation, TR-653-NCQA, 2009). In this context, reliability represents the ability of a measure to confidently distinguish the performance of one physician from another. As discussed in the report: “Conceptually, it is the ratio of signal to noise. The signal in this case is the proportion of variability in measured performance that can be explained by real differences in performance. There are 3 main drivers of reliability; sample size, differences between physicians, and measurement error.”

According to this approach, reliability is estimated with a beta-binomial model. The beta-binomial model is appropriate for measuring the reliability of pass/fail measures such as those proposed.

Physician specific reliability is around .9 for each year, and thus can be considered to be good. Reliability scores vary from 0.0 to 1.0, with a score of zero indicating that all variation is attributable to measurement error (noise, or variation across patients within providers) whereas a reliability of 1.0 implies that all variation is caused by real difference in performance across accountable entities. There is not a clear cut-off for minimum reliability level. Values of 0.7, however, are considered sufficient to see differences between some physicians and the mean (see RAND tutorial, 2009). The Results of the Signal to Noise analysis can be found in Table 3 for Timing to Surgery.

Table 3. Reliability Statistics from the Signal to Noise Analysis: Timing to Surgery

Year	# of Physicians	Reliability Statistic from signal-to-noise analysis (95% CI)
2012	69	.88 (.85,.91)
2013	69	.89 (.85,.92)

2014	59	.91 (.88,.95)
2015	59	.91 (.88,.93)
2016	52	.90 (.87,.93)

Performance Gap

This candidate measure shows a moderately high overall computed compliance rate in our sample dataset, derived from the Cleveland Clinic with individual physicians’ means ranging from .89 to .94 for the years 2012-2016. Distributions can be found in table 4b.1.

In order to address the possibility for variance in care within that overall mean compliance rates, this group performed a secondary analysis, evaluating only practitioners who were not 100% compliant (4b.2). Within this secondary analysis, the gap increases by about 10% on average and the means range from .75 in 2012 to .81 in 2014. This secondary analysis combined with the expectation that the analyzed data were obtained from a high-quality academic research hospital with a compliance rate likely higher than the national norm lead this group to believe a reasonable performance gap exists.

Table 4b.1 Minimum to Maximum Ranges of Performances scores for All and Compliant Physicians

Year	Mean	SD	Max	99%	95%	90%	75%Q3	50% Median	25% Q1	10%	5%	1%	Min
2012	0.89483	0.198898	1	1	1	1	1	1	0.857143	0.7	0.666667	0	0
2013	0.901951	0.163173	1	1	1	1	1	1	0.830508	0.695652	0.666667	0	0
2014	0.939034	0.102608	1	1	1	1	1	1	0.888889	0.75	0.722222	0.666667	0.666667
2015	0.914594	0.166035	1	1	1	1	1	1	0.875	0.75	0.6	0	0
2016	0.885474	0.186951	1	1	1	1	1	0.98	0.828431	0.75	0.5	0	0

Table 4b.2 Minimum to Maximum Ranges of Performances scores for All and Compliant Physicians

Year	Mean	SD	Max	99%	95%	90%	75%Q3	50% Median	25% Q1	10%	5%	1%	Min
2012	0.749768	0.241827	0.947368	0.947368	0.941176	0.933333	0.875	0.823529	0.714286	0.333333	0	0	0
2013	0.758379	0.176302	0.923077	0.923077	0.9	0.888889	0.870833	0.8	0.722826	0.666667	0.5	0	0
2014	0.810685	0.090976	0.96	0.96	0.96	0.928571	0.888889	0.777778	0.75	0.666667	0.666667	0.666667	0.666667
2015	0.770955	0.20414	0.923077	0.923077	0.923077	0.916667	0.885714	0.833333	0.75	0.6	0.5	0	0
2016	0.770948	0.209804	0.96	0.96	0.95	0.947368	0.92	0.828431	0.75	0.5	0.5	0	0

References

1. Adams, John L. "The Reliability of Provider Profiling: A Tutorial." RAND Corporation. N.p., 25 June 2009. Web. 11 May 2017.
2. "Guidance for Measure Testing and Evaluating Scientific Acceptability of Measure Properties." NQF: Measure Testing Task Force. N.p., n.d. Web. 07 Dec. 2016. Available at: http://www.qualityforum.org/Publications/2011/01/Measure_Testing_Task_Force.aspx
3. Elliott J, Beringer T, Kee F, Marsh D, Willis C, Stevenson M. Predicting survival after treatment for fracture of the proximal femur and the effect of delays to surgery. *J Clin Epidemiol* 2003;56(8):788-795.
4. Fox HJ, Pooler J, Prothero D, Bannister GC. Factors affecting the outcome after proximal femoral fractures. *Injury* 1994;25(5):297-300.
5. McGuire KJ, Bernstein J, Polsky D, Silber JH. The 2004 Marshall Urist award: delays until surgery after hip fracture increases mortality. *Clin Orthop Relat Res* 2004;(428):294-301.
6. Moran CG, Wenn RT, Sikand M, Taylor AM. Early mortality after hip fracture: is delay before surgery important? *J Bone Joint Surg Am* 2005;87(3):483-489.
7. Novack V, Jotkowitz A, Etzion O, Porath A. Does delay in surgery after hip fracture lead to worse outcomes? A multicenter survey. *Int J Qual Health Care* 2007;19(3):170-176.
8. Orosz GM, Magaziner J, Hannan EL et al. Association of timing of surgery for hip fracture and patient outcomes. *JAMA* 2004;291(14):1738-1743.
9. Parker MJ, Pryor GA. The timing of surgery for proximal femoral fractures. *J Bone Joint Surg Br* 1992;74(2):203-205.
10. Radcliff TA, Henderson WG, Stoner TJ, Khuri SF, Dohm M, Hutt E. Patient risk factors, operative care, and outcomes among older community-dwelling male veterans with hip fracture. *J Bone Joint Surg Am* 2008;90(1):34-42.
11. Siegmeth AW, Gurusamy K, Parker MJ. Delay to surgery prolongs hospital stay in patients with fractures of the proximal femur. *J Bone Joint Surg Br* 2005;87(8):1123-1126.

Disclaimer:

These Performance Measures and related data specifications were developed by the American Academy of Orthopaedic Surgeons (AAOS) through a multi-disciplinary physician workgroup and are based on a systematic review of published literature and/or relevant clinical practice guidelines to facilitate quality improvement activities by physicians. These Performance Measures are not clinical guidelines and do not establish a standard of medical care, and have not been tested for all potential applications. They are not intended to establish fixed protocols, but rather to serve as metrics by which a health care provider's or facility's performance may be compared with national benchmarks. Patient care and treatment should always be based on the clinician's independent medical judgment, given the individual patient's clinical circumstances. The Performance Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, for example, use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Performance Measures require a license agreement between the user and the AAOS. The AAOS nor its members shall be responsible for any use of the Performance Measures.

Appendix A. Disclosures

William Timothy Brox, MD (This individual reported nothing to disclose); Submitted on: 01/28/2015

David B Carmack, MD Submitted on: 12/20/2014

OTA (I served on the OTA Health Policy Committee as a non-paid volunteer): Board or committee member (\$0)

Charles M Davis III, MD Submitted on: 12/22/2014

AAOS: Board or committee member (\$0) (Self)

American Association of Hip and Knee Surgeons: Board or committee member (\$0) (Self)

Journal of Arthroplasty: Editorial or governing board (\$0) (Self)

Eitan Dickman, MD Submitted on: 12/20/2014

Society for Academic Emergency Medicine Emergency Ultrasound Academy BOD: Board or committee member (\$0) This is an unpaid position(Self)

Stephen L Kates, MD Submitted on: 10/26/2014

Sage Publications: Editorial or governing board (\$0)

AAOS: Board or committee member (\$0)

AO Foundation: Paid presenter or speaker (\$0) Number of Presentations: 0

AO North America: Board or committee member (\$0)

AOTrauma: Board or committee member (\$0)

Orthopaedic Trauma Association: Board or committee member (\$0)

Sage Publications: Publishing royalties, financial or material support (\$0)

Surgical Excellence: Paid consultant (\$0)

Synthes: Research support (\$0)

James F Kellam, MD Submitted on: 12/19/2014

AO Foundation: Editorial or governing board (\$0)

AONA: Editorial or governing board (\$0)

Orthopaedic Trauma Association: Board or committee member (\$0)

Douglas W Lundy, MD Submitted on: 11/18/2014

AAOS: Board or committee member (\$0) (Self) - chair Medical Liability Committee, member Council on Advocacy

American Board of Orthopaedic Surgery, Inc.: Board or committee member (\$0) Board of Directors

American College of Surgeons: Board or committee member (\$0)

Clinical Orthopaedics and Related Research: Editorial or governing board (\$0) (Self) - journal reviewer

Georgia Orthopaedic Society: Board or committee member (\$0) (Self) - Board of Directors

Journal of Orthopaedic Trauma: Editorial or governing board (\$0) Associate editor(Self)

Journal of the Southern Medical Association: Editorial or governing board (\$0) Reviewer(Self)

Livengood Engineering: Stock or stock Options Number of Shares: 100 (Self)

Orthopaedic Trauma Association: Board or committee member (\$0) (Self) - Member-at-Large - Board of

Directors Orthopedics: Editorial or governing board (\$0) (Self) - journal reviewer

Kevin Means, MD Submitted on: 12/26/2014

American Academy of Physical Medicine and Rehabilitation: Board or committee member (\$0)

Membership Committee(Self)

Association of Academic Physiatrists: Board or committee member (\$1,100) Rehabilitation Medicine Scientist Training Program Advisory Board(Self)
Demos Medical Publishing: Publishing royalties, financial or material support (\$600) Royalties for editing a Geriatrics book in 2013 (Self)

Simon Mears, MD Submitted on: 12/03/2014

International Geriatric Fracture Society: Board or committee member (\$0)
Journal of Geriatric Orthopaedics and Rehabilitation Arthritis and Rheumatism EPlasty: Editorial or governing board(\$0)
Journal of the American Geriatrics Society: Editorial or governing board (\$0)

Daniel Ari Mendelson, MD, MS, FACP, AGSF Submitted on: 01/29/2015

Geriatric Orthopaedic Surgery and Rehabilitation/Sage: Editorial or governing board (\$0) (Self) Editorial Board Member

Anna Noel Miller, MD Submitted on: 01/29/2015

DePuy, A Johnson & Johnson Company: Paid consultant (\$500) N/A(Self)

Arvind D Nana, MD Submitted on: 02/02/2014

AAOS: Board or committee member; Board or committee member (\$0) (Self) Liaison to the National Bone Health Alliance
Advanced Orthopaedic Solutions: Unpaid consultant Unpaid consultant(Self)
American Orthopaedic Association: Board or committee member (\$0) (Self) Own the Bone Multi-Disciplinary Advisory Board
International Geriatric Fracture Society: Board or committee member (\$0)
Musculoskeletal Infection Society: Board or committee member (\$0) (Self) Past President
Orthopaedic Trauma Association: Board or committee member (\$0) (Self) Evidence Based Value & Quality Committee

Laura Lowe Tosi, MD Submitted on: 01/02/2015

American Orthopaedic Association: Society for Women's Health Research: Board or committee member (\$0)
Bone and Joint Decade, U.S.A.: Board or committee member (\$0)
Journal of Children's Orthopaedics: Editorial or governing board (\$0)
Medical Society of the District of Columbia: Board or committee member (\$0)
National Bone Health Alliance: Board or committee member (\$0)
Osteogenesis Imperfecta Foundation: Board or committee member (\$0)

Colleen Walsh, NP (This individual reported nothing to disclose); Submitted on: 12/11/2014

Douglas M White, DPT Submitted on: 01/23/2015

American Physical Therapy Assoc Orthopaedic Section: Board or committee member (\$0) Imaging Special Interest Group, Clinical Practice Guidelines Hip Panel(Self)

Douglas G Wright, MD (This individual reported nothing to disclose); Submitted on: 01/07/2015