

## Table 2 Demographic Data on Revision Total Hip Arthroplasty According to United States Census Region

Code	Northeast	Midwest	South	West	Total
00.70: All-component revision	3,445 (16.3%)	5,090 (24.1%)	7,582 (35.9%)	4,988 (23.6%)	21,105 (100.0%)
00.71: Acetabular component revision	997 (15.3%)	1,638 (25.1%)	2,329 (35.6%)	1,572 (24.0%)	6,537 (100.0%)
00.72: Femoral component revision	1,231 (18.2%)	1,777 (26.3%)	2,433 (36.0%)	1,309 (19.4%)	6,749 (100.0%)
00.73: Isolated head/liner revision	1,160 (17.9%)	1,358 (21.0%)	2,191 (33.8%)	1,770 (27.3%)	6,479 (100.0%)
80.05: Arthrotomy/removal	697 (14.9%)	1,111 (23.7%)	1,971 (42.1%)	903 (19.3%)	4,682 (100.0%)
81.53: Hip revision, not otherwise specified	1,011 (20.7%)	1,050 (21.5%)	2,078 (42.6%)	742 (15.2%)	4,880 (100.0%)
Other combinations	188 (20.6%)	192 (21.0%)	282 (30.9%)	252 (27.6%)	913 (100.0%)
<b>Total</b>	<b>8,729 (17.0%)</b>	<b>12,214 (23.8%)</b>	<b>18,867 (36.7%)</b>	<b>11,535 (22.5%)</b>	<b>51,345 (100.0%)</b>

Reprinted with permission from Bozic KJ, Kurtz SM, Lau E, Ong K, Vail TP, Berry DJ: The Epidemiology of Revision Total Hip Arthroplasty in the United States. J Bone Joint Surg Am 2009;91:128-133.